

BARBICIDE®

Hospital

GERMICIDE, PSEUDOMONACIDE FUNGICIDE & VIRUCIDE*

Anti Rust Formula

ACTIVE INGREDIENT:	
Alkyl (50% C ₁₀ , 40% C ₁₂ , 10% C ₁₆) dimethyl benzyl ammonium chloride	5.12%
INERT INGREDIENTS	94.88%
TOTAL:	100.00%
(Contains sodium nitrite)	

KEEP OUT OF REACH OF CHILDREN DANGER

Call poison control center or doctor immediately for treatment advice. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. IF SWALLOWED: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents 64 FL. OZ. 1.89 L

KILLS *HIV-1 (AIDS VIRUS), *HBV AND *HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD AND BODY FLUIDS.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS.

On pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

PERSONAL PROTECTION

When handling items soiled with blood or body fluids, use appropriate barrier protection such as disposable latex gloves, gowns, masks and eye coverings.

CLEANING PROCEDURE

Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant.

DISPOSAL OF INFECTIOUS MATERIALS

Blood and other body fluids must be autoclaved and disposed of according to Federal, State and local regulations for infectious waste disposal.

CONTACT TIME

Leave surface wet for 10 minutes.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: KEEP OUT OF REACH OF CHILDREN. CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE AND SKIN BURNS. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD. WEAR EYE PROTECTION (FACE SHIELD AND/OR GOGGLES), PROTECTIVE CLOTHING, AND RUBBER GLOVES WHEN HANDLING. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING AND BEFORE, DRINKING, OR USING TOBACCO. REMOVE CONTAMINATED CLOTHING AND WASH BEFORE REUSE.

STORAGE AND DISPOSAL

Do not store on side. Avoid creasing or impacting of side walls. Non-refillable container. Do no reuse or refill this container. Offer for recycling, if available or wrap container and put in trash.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinse water is a violation of Federal Law. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

A PRODUCT OF
KING
RESEARCH®

7025 West Marcia Road
MILWAUKEE, WI 53223

EPA Registration No. 954-11 EPA Establishment No. 051435-WI-004



BARBICIDE 64 oz.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. To make a HOSPITAL DISINFECTANT mix 2 ounces of BARBICIDE in 32 ounces of water. BARBICIDE is for use in hospitals, health care facilities, tanning facilities and in beauty salons, barbershops, and hairstyling establishments. SURFACES: On inanimate, hard non-porous surfaces such as floors, sink tops, tables, acrylic tanning beds, telephones, toilet seats, and door handles. Apply solution prepared as above to pre-cleaned surfaces with a cloth, mop, sponge, or coarse mist pump or trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes and then let air-dry. BEAUTY/BARBER INSTRUMENTS AND TOOLS: Thoroughly clean brushes, combs, clipper blades, shears, razors, and other inanimate hard surfaced non-porous tools and implements prior to complete immersion for 10 minutes (or as required by local authorities) in a solution as prepared above. Rinse to remove BARBICIDE after immersion and dry before use. Fresh solution must be prepared daily or more often if solution becomes diluted or soiled. PERFORMANCE: When tested with full immersion for 10 minutes in a solution prepared as above, BARBICIDE has been found to be effective against: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus (including Staphylococcus aureus Methicillin Resistant (MRSA)), Trichophyton mentagrophytes, *Herpes Simplex Virus Type-1 and Type-2, *Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS virus), Influenza A Virus (in 5% Serum), *Hepatitis B Virus and *Hepatitis C Virus.
BAR-1890-01